

General

Title

Stroke and stroke rehabilitation: percentage of patients aged 18 years and older with the diagnosis of *ischemic stroke* or *TIA* who were prescribed antiplatelet therapy at discharge.

Source(s)

American Academy of Neurology, American College of Radiology, Physician Consortium for Performance Improvement \hat{A} \mathbb{R} , National Committee for Quality Assurance. Stroke and stroke rehabilitation physician performance measurement set. Chicago (IL): American Medical Association (AMA), National Committee for Quality Assurance (NCQA); 2009 Feb. 20 p.

Measure Domain

Primary Measure Domain

Process

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

Secondary Measure Domain

Does not apply to this measure

Brief Abstract

Description

This measure is used to assess the percentage of patients aged 18 years and older with the diagnosis of ischemic stroke OR transient ischemic attack (TIA) who were prescribed antiplatelet therapy at discharge.

To promote a comprehensive approach to performance improvement, the Stroke and Stroke Rehabilitation Measurement Set is intended for use in its entirety when measuring clinical quality in the care of eligible patients. Full implementation of this measurement set for patients with a diagnosis of ischemic stroke or transient ischemic attack (TIA) should always include the Physician Consortium for Performance Improvement® measure, Radiology: Stenosis Measurement in Carotid Imaging Reports.

Rationale

Following a stroke, patients should be on antiplatelet therapy to decrease the risk of additional strokes.*

*The following clinical recommendation statements are quoted <u>verbatim</u> from the referenced clinical guidelines and represent the evidence base for the measure:

We recommend that every patient who has experienced a noncardioembolic (atherothrombotic, lacunar, or cryptogenic) stroke or transient ischemic attack (TIA) and has no contraindication receives an antiplatelet agent regularly to reduce the risk of recurrent stroke and other vascular events. Aspirin, 50 to 325 mg qd; the combination of aspirin, 25 mg, and extended-release dipyridamole, 200 mg bid; or clopidogrel, 75 mg qd, are all acceptable options for initial therapy. (American College of Chest Physicians [ACCP]) (Grade 1A)

For patients with noncardioembolic ischemic stroke or TIA, antiplatelet agents rather than oral anticoagulation are recommended to reduce the risk of recurrent stroke and other cardiovascular events. (American Stroke Association [ASA])

Aspirin (50 to 325 mg/d), the combination of aspirin and extended-release dipyridamole, and clopidogrel are all acceptable options for initial therapy. (ASA)

Primary Clinical Component

Ischemic stroke; transient ischemic attack (TIA); antiplatelet therapy

Denominator Description

All patients aged 18 years and older with the diagnosis of ischemic stroke OR transient ischemic attack (TIA) (see the related "Denominator Inclusions/Exclusions" field)

Numerator Description

Patients who were prescribed antiplatelet therapy at discharge

Evidence Supporting the Measure

Evidence Supporting the Criterion of Quality

A clinical practice guideline or other peer-reviewed synthesis of the clinical evidence

Evidence Supporting Need for the Measure

Need for the Measure

Use of this measure to improve performance

Evidence Supporting Need for the Measure

McGlynn EA, Asch SM, Adams J, Keesey J, Hicks J, DeCristofaro A, Kerr EA. The quality of health care delivered to adults in the United States. N Engl J Med. 2003 Jun 26;348(26):2635-45. PubMed

Reeves MJ, Arora S, Broderick JP, Frankel M, Heinrich JP, Hickenbottom S, Karp H, LaBresh KA, Malarcher A, Mensah G, Moomaw CJ, Schwamm L, Weiss P, Paul Coverdell Prototype Registries Writing Group. Acute stroke care in the US: results from 4 pilot prototypes of the Paul Coverdell National Acute Stroke Registry. Stroke. 2005 Jun;36(6):1232-40. PubMed

Thom T, Haase N, Rosamond W, Howard VJ, Rumsfeld J, Manolio T, Zheng ZJ, Flegal K, O'Donnell C, Kittner S, Lloyd-Jones D, Goff DC Jr, Hong Y, Adams R, Friday G, Furie K, Gorelick P, Kissela B, Marler J, Meigs J, Roger V, Sidney S, Sorlie P, Steinberger J, Wasserthiel-Smoller S, Wilson M, Wolf P. Heart disease and stroke statistics-2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Circulation. 2006 Feb 14;113(6):e85-151. PubMed

State of Use of the Measure

State of Use

Current routine use

Current Use

Internal quality improvement

National reporting

Application of Measure in its Current Use

Care Setting

Hospitals

Professionals Responsible for Health Care

Advanced Practice Nurses

Physician Assistants

Physicians

Lowest Level of Health Care Delivery Addressed

Individual Clinicians

Target Population Age

Age greater than or equal to 18 years

Target Population Gender

Either male or female

Stratification by Vulnerable Populations

Unspecified

Characteristics of the Primary Clinical Component

Incidence/Prevalence

Unspecified

Association with Vulnerable Populations

Unspecified

Burden of Illness

Unspecified

Utilization

Unspecified

Costs

Unspecified

Institute of Medicine (IOM) Healthcare Quality Report Categories

IOM Care Need

Staying Healthy

IOM Domain

Effectiveness

Data Collection for the Measure

Case Finding

Users of care only

Description of Case Finding

All patients aged 18 years and older with the diagnosis of ischemic stroke OR transient ischemic attack (TIA)

Denominator Sampling Frame

Patients associated with provider

Denominator Inclusions/Exclusions

Inclusions

All patients aged 18 years and older with the diagnosis of ischemic stroke OR transient ischemic attack (TIA)

Exclusions

Documentation of medical reason(s) for not prescribing antiplatelet therapy at discharge (including identification from medical record that the patient is on anticoagulation therapy)

Documentation of patient reason(s) for not prescribing anticoagulation therapy at discharge

Relationship of Denominator to Numerator

All cases in the denominator are equally eligible to appear in the numerator

Denominator (Index) Event

Clinical Condition

Institutionalization

Denominator Time Window

Time window brackets index event

Numerator Inclusions/Exclusions

Inclusions

Patients who were prescribed antiplatelet therapy at discharge

Exclusions

None

Measure Results Under Control of Health Care Professionals, Organizations and/or Policymakers

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

Numerator Time Window

Institutionalization

Data Source

Administrative data

Level of Determination of Quality

Individual Case

Pre-existing Instrument Used

Unspecified

Computation of the Measure

Scoring

Rate

Interpretation of Score

Better quality is associated with a higher score

Allowance for Patient Factors

Unspecified

Standard of Comparison

Internal time comparison

Evaluation of Measure Properties

Extent of Measure Testing

Unspecified

Identifying Information

Original Title

Measure #2: discharged on antiplatelet therapy.

Measure Collection Name

The Physician Consortium for Performance Improvement® Measurement Sets

Measure Set Name

Stroke and Stroke Rehabilitation Physician Performance Measurement Set

Submitter

American Medical Association on behalf of the American Academy of Neurology, American College of Radiology, Physician Consortium for Performance Improvement®, and National Committee for Quality Assurance - Medical Specialty Society

Developer

American Academy of Neurology - Medical Specialty Society

American College of Radiology - Medical Specialty Society

National Committee for Quality Assurance - Health Care Accreditation Organization

Physician Consortium for Performance Improvement® - Clinical Specialty Collaboration

Funding Source(s)

Unspecified

Composition of the Group that Developed the Measure

Joseph Drozda, Jr., MD (*Co-Chair*); Robert Holloway, MD, MPH (*Co-Chair*); David Seidenwurm, MD (*Co-Chair*); Chris Alexander, III, MD, FACP; John Barr, MD; Oscar Benavente, MD, FRCPC; Christopher Bever, Jr., MD, MBA; Thomas Bleck, MD FCCM; Ronald Gabel, MD; Judith Hinchey, MD; Irene Katzan, MD; Rahul K. Khare, MD; Michael Lev, MD; John McMahon, MD; Mark Morasch, MD, FACS; Suresh Mukherji, MD; Konrad C. Nau, MD; Sam J. W. Romeo, MD, MBA; Eric Russell, MD, FACR; Marilyn Rymer, MD; Ajay Sharma, MD; John Schneider, MD, PhD; Timothy Shephard, PhD; Michael A. Sloan, MD, MS; William D. Smucker, MD, CMD; Patrick Turski, MD, FACR; Linda Williams, MD; Richard Zorowitz, MD

Sarah Tonn, MPH, American Academy of Neurology

Sandra H. Bjork, RN, JD, American College of Radiology

Penelope Solis, JD, American Heart Association/American Stroke Association

Andrea Gelzer, MD, MS, FACP, Health Plan Representative

Mark Antman, DDS, MBA, American Medical Association; Karen S. Kmetik, PhD, American Medical Association; Beth Tapper, MA, American Medical Association; Samantha Tierney, MPH, American Medical Association

Rebecca A. Kresowik, Facilitator; Timothy F. Kresowik, MD, Facilitator

Financial Disclosures/Other Potential Conflicts of Interest

Conflicts, if any, are disclosed in accordance with the Physician Consortium for Performance Improvement® conflict of interest policy.

Endorser

Included in

Ambulatory Care Quality Alliance

Physician Quality Reporting Initiative

Adaptation

This measure was harmonized to the extent possible with hospital level measures for stroke developed by The Joint Commission.

Parent Measure

Unspecified

Release Date

2006 Sep

Revision Date

2009 Feb

Measure Status

This is the current release of the measure.

The Physician Consortium for Performance Improvement reaffirmed the currency of this measure in November 2010.

Source(s)

American Academy of Neurology, American College of Radiology, Physician Consortium for Performance Improvement®, National Committee for Quality Assurance. Stroke and stroke rehabilitation physician performance measurement set. Chicago (IL): American Medical Association (AMA), National Committee for Quality Assurance (NCQA); 2009 Feb. 20 p.

Measure Availability

The individual measure, "Measure #2: Discharged on Antiplatelet Therapy," is published in the "Stroke and Stroke Rehabilitation: Physician Performance Measurement Set." This document and technical specifications are available in Portable Document Format (PDF) from the American Medical Association (AMA)-convened Physician Consortium for Performance Improvement® Web site:

10/10/10/	nhysicianconsortium ord

For further information, please contact AMA staff by e-mail at cqi@ama-assn.org.

NQMC Status

This NQMC summary was completed by ECRI Institute on September 13, 2007. The information was verified by the measure developer on October 26, 2007. This NQMC summary was edited by ECRI Institute on September 28, 2009. The information was reaffirmed by the measure developer on November 17, 2010.

Copyright Statement

- © Measures including specifications
- © 2005-6 American Medical Association and National Committee for Quality Assurance

Disclaimer

NQMC Disclaimer

The National Quality Measures Clearinghouseâ, ¢ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. Moreover, the views and opinions of developers or authors of measures represented on this site do not necessarily state or reflect those of NQMC, AHRQ, or its contractor, ECRI Institute, and inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.